


SYSTEMATIC MAP PROTOCOL

Open Access



Collating existing evidence on cumulative impacts of invasive plant species in riparian ecosystems of British Columbia, Canada: a systematic map protocol

Fabio Mologni^{1*} , Chandra E. Moffat^{2,1} and Jason Pither¹

Abstract

Background Globally, the structure and functioning of foreshore and riparian ecosystems are being dramatically impacted by non-native invasive plant species. Invasive species can outcompete and replace native species, modify geochemical and hydraulic cycles, alter trophic processes, and change the composition and structure of communities above and below ground. However, these impacts are often investigated in isolation, even though one invasive species might increase or mitigate the impacts of others (i.e. cumulative impacts), potentially with cascading effects. Although cumulative impacts have long been studied within other environmental contexts, research on the cumulative impacts of invasive species is comparatively scarce. We aim to develop a protocol to systematically identify and collate evidence on the individual and cumulative impacts of a set of plant species invasive in foreshore and riparian ecosystems of British Columbia, Canada. Our primary question is: what evidence is available on the individual and cumulative impacts of invasive plants in the riparian and foreshore ecosystems of British Columbia, Canada? In addition, our systematic map will identify the strengths and gaps in knowledge pertaining to invasive plant species impacts in foreshore and riparian ecosystems, with the ultimate goal of facilitating the development of evidence-based management strategies.

Methods We identified the research topic and the primary and secondary questions with the support of stakeholders. We then devised a flexible string that allows for searching target invasive species. Using this string, we searched the literature for pilot species that aided the iterative development of the protocol. Once all target species are identified, we will carry out a systematic literature search on their impacts. We will search Web of Science and the CABI compendium for invasive species. We will include studies if they (i) refer to the target invasive species, (ii) focus on its environmental impacts and (iii) investigate such impacts in riparian ecosystems (iv) within North America (i.e. Canada and U.S.A.). We will use a two-stage screening process: titles and abstracts first, then the full manuscript. From each source, we will extract impact description, ecosystem component impacted, and magnitude and directionality of impacts. We will include a publicly available database of studies, descriptive statistics, and a narrative summary within our synthesis outcomes.

Keywords Cumulative impacts, British Columbia, Invasive species, Impacts, Riparian ecosystems, Plant invasions, Foreshore ecosystems, Protocol, Systematic maps

*Correspondence:

Fabio Mologni

fabio.mologni@ubc.ca

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Biological invasions in foreshore and riparian ecosystems

Foreshore and riparian ecosystems are vitally important from ecological, cultural, and economic standpoints. Although their spatial extent is small, they are often hotspots of biodiversity, hosting rare species, and serving as refugia and corridors essential to many others [1–3]. These ecosystems also provide essential functions and services such as improving water quality, flood mitigation, and minimizing erosion [2, 4, 5]. As such, foreshore and riparian habitats are the focus of targeted management and conservation strategies in many countries [6–9].

Despite their recognized importance, foreshore and riparian ecosystems are being impacted by many anthropogenic stressors [10]. Infrastructures (e.g. dams, dyking, channelization) and water management (e.g. water diversion, irrigation, dredging) can radically modify water levels and flow and disrupt natural fluvial dynamics [1, 5, 11, 12]. Contamination and nutrient additions can alter water quality, reduce biodiversity, and promote bioaccumulation [1, 13]. Habitat loss through agriculture, deforestation, and development disproportionately impacts foreshore and riparian zones [1, 14–16], and was estimated to be up to two-thirds in the U.S. alone [17]. Additionally, freshwater ecosystems are oftentimes highly invaded by non-native species due to their proximity to human settlements and their function as dispersal corridors [14, 18–21].

Invasive species can impact riparian ecosystems in various ways, but invasive plants have particularly pervasive impacts on ecosystem structure and functioning. By spreading aggressively, they displace both plant and animal native species [22–25], modify geochemical and hydraulic cycles [26, 27], alter trophic processes [28], and change the composition and structure of communities above and below ground [2, 29]. Additionally, invasive plants alter traditional practices and resource use by Indigenous peoples [28]. The cumulative impacts of invasive plants on riparian ecosystems are potentially profound, but research to quantify such effects remains limited [2, 30].

Here, we aim to develop a framework for systematically collating and mapping evidence on the individual and cumulative impacts of plant species that are invasive within foreshore and riparian ecosystems, and we will apply our protocol to systems in British Columbia, Canada.

Individual and cumulative impacts: definitions, examples and previous work

In invasion ecology, individual impacts are defined as measurable changes caused by non-native species on a target ecosystem [31, 32]. They can vary greatly in type, magnitude, and directionality. For instance, some impacts might be barely detectable (e.g. gene flow through hybridization), while others can produce pronounced, observable effects (e.g. ecosystem dominance). Impacts can be direct (e.g. displacement of native species), but also mediated through other factors (e.g. competition for resources, [31]). Finally, while non-native species have been investigated in large part because of their negative effects, impacts can vary along a continuum from negative to positive [32, 33], and can be ecosystem or context-dependent.

Identifying an impact's directionality presents some challenges. Negative impacts are typically equated to unfavourable outcomes for humans [32]. However, this approach is strongly biased by the value system and worldview of the researcher [33, 34]. In an effort to minimize subjectivity and value-based identifications of impact directionality, we define as negative or positive any quantifiable reduction or increase in ecosystem properties or attributes (e.g. native species richness and abundance, nutrient cycling, water quality, etc., [32]). For instance, we define as positive an increase in the fitness or number of individuals of a native species but as negative its reduction.

The combination and interaction of multiple individual impacts are referred to as cumulative impacts and many definitions of this concept exist. For the Canadian Environmental Assessment Act (CEAA), they are “*changes to the environment that are caused by an action in combination with other past, present and future human actions*” [35]. The Council on Environmental Quality (CEQ) suggests impacts have to be incremental [36]. The most well-articulated definition is that of the European Environmental Agency (EEA), which defines them as: “*the impacts (positive or negative, direct and indirect, long-term and short-term impacts) arising from a range of activities throughout an area or region, where each individual effect may not be significant if taken in isolation. Such impacts can arise from the growing volume of traffic, the combined effect of a number of agriculture measures leading to more intensive production and use of chemicals, etc. Cumulative impacts include a time dimension, since they should calculate the impact on environmental resources resulting from changes brought about by past, present and reasonably foreseeable future actions.*” [37]. Consistent elements among these definitions are (1) the combination of multiple individual impacts, (2) a time component and (3) the human agency. While not

explicitly stated in the previous definitions, cumulative impacts also have a spatial dimension, or they can accumulate in space as well as temporally [38].

We define cumulative impacts in biological invasions as the combined effect of multiple impacts when at least one is generated by an invasive species. Cumulative impacts include recurrent impacts of a single species and the combined effect of multiple invaders, but also the compounded impact of invading species and other anthropogenic stressors [12]. Our definition incorporates all the elements of previous definitions; however, it is more restrictive, as the primary focus is the impacts of invasive species. Conversely, it includes impacts of any magnitude, type or directionality.

The term ‘cumulative’ might imply that the total effect of multiple impacts is always greater than that of individual impacts. Multiple invaders can collectively increase native species displacement, or enhance topsoil nutrient concentration (additive impacts, [29, 39]). An N-fixer might increase soil nitrogen, facilitating invasions by more competitive nitrophilous species, which in turn will displace natives (multiplicative impacts, [29]). However, additive or multiplicative impacts are not the only potential outcomes. Competition between two invaders might instead reduce their impact per capita. For example, an allopathic species might negatively affect both native and non-native species. In this case, one invader mitigates the impacts of another invader [38].

Despite a long history of research on cumulative impacts within environmental contexts [38], the literature on the cumulative impacts of invasive species is relatively scarce. Most work in biological invasions focuses on a single species or single direct impact [40–45]. Even when multiple impacts are identified, their cumulative effect is rarely considered [30, 46]. This is despite previously proposed theoretical frameworks share some conceptual overlap. One such example is the invasion meltdown, which posits that interactions among invaders might increase their impacts [47]. Critically for our work, little research effort explored the cumulative impacts of invasive plant species in riparian and foreshore ecosystems. Therefore, anticipating a lack of studies on cumulative impacts, we will also include individual impacts in this systematic map.

Topic identification and stakeholder input

There is a clear need for work identifying the cumulative impacts of invasive species in riparian ecosystems. The Province of British Columbia, Ministry of Forests Invasive Plant Program, highlighted the need to synthesize current evidence on the impacts of invasive plant species in riparian and foreshore ecosystems within the province, to inform research and management needs.

British Columbia’s riparian and foreshore ecosystems are invaded by numerous highly destructive invasive plant species, such as Russian Olive (*Elaeagnus angustifolia*), Phragmites (*Phragmites australis*), Knotweeds (*Reynoutria* spp., syn. *Fallopia*), Tree of Heaven (*Ailanthus altissima*) and Canary reed grass (*Phalaris arundinacea*). While the impacts of these species have been extensively investigated [42, 48–52], there is no comprehensive assessment of their cumulative impacts.

Stakeholders in the provincial government played a pivotal role in shaping the research topic and refining the scope of the systematic map. Stakeholders include the British Columbia Ministry of Forests, Agriculture and Agri-Food Canada, and the University of British Columbia. Based on their expert knowledge and the available data, they provided a list of 10–15 plant species that are invasive in the target ecosystems and geographic areas, thereby aiding in the identification of specific research questions and objectives. Input from practitioners and other researchers helped refine the approach and the methodology. Through ongoing dialogue and feedback, stakeholders were able to establish clear expectations, develop a robust methodology, and identify appropriate outcomes for the systematic map. In addition to quantifying the cumulative impacts of plant species invasive to riparian ecosystems, stakeholders have identified two additional aspects as essential. First is the development of a reproducible protocol that can be employed in future systematic studies of invasive species impacts. Second is the investigation of how the cumulative impacts of invasive species will vary under current climate change scenarios.

Protocols are a crucial aspect of developing a project, particularly in the case of systematic work [53]. Good protocols need to be transparent, detailed and reproducible, allowing other researchers to replicate their work [53–56]. In this case, we do not simply want to describe our procedure for mapping the existing literature, but we specifically aim to provide a tool that is sufficiently flexible and reproducible to be applied in the investigation of other invasive species or ecosystems.

Climate change is a key contributor to the cumulative impacts of invasive species across both terrestrial and aquatic ecosystems. However, the nature and magnitude of its effect of invasive species’ impacts is often unclear. Interactions between particular invasive plants and the diverse facets of climate change are challenging to predict and likely species- and context-dependent [57]. For instance, while the ranges of many non-native invasive species may expand as temperature rises [58], others may contract or shift in response to both abiotic and biotic factors [57, 59]. Nevertheless, strategies for mitigating negative impacts are sorely needed. A key first step is

synthesizing the diverse and extensive research on this topic.

Here, we present a reproducible systematic map protocol [53] for screening, collating, and describing research on the impacts of priority invasive plants in riparian and foreshore ecosystems, and we will apply it to systems in British Columbia. Given their efficacy and comprehensiveness, systematic maps are increasingly common in environmental management [54]. Through the systematic map process, we will identify knowledge clusters and gaps (i.e. areas of high and low concentration of the research effort), and synthesize results within the context of current climate change scenarios. Key outputs will include (1) a robust analytical framework for qualitatively predicting—based on the best available evidence—the cumulative impacts of invasive plants under changing climates and followed by (2) a more detailed assessment for a selection of priority invasive plant species (identified by the BC Ministry of Forests Invasive Alien Plant Program). These outputs will have high utility for policy, planning and strategic, evidence-based decision management of ecosystems impacted by priority invasive plant species in British Columbia.

Objective of the review

We aim to systematically collate and map evidence on the individual and cumulative impacts of a selection of plant species invasive to riparian ecosystems in British Columbia, Canada.

Primary question

What evidence is available on the individual and cumulative impacts of invasive plants in the riparian and foreshore ecosystems of British Columbia, Canada?

Components of the primary question

- Population: Riparian and foreshore ecosystems in British Columbia
- Exposure: non-native plant species invasive to riparian and foreshore ecosystems of British Columbia
- Comparator: No impact or absence of invasive plant species.
- Outcome: A synthesis of both the individual and collective cumulative impacts of the selected invasive plant species

Secondary question

We will describe variations in the research effort with regard to:

- Geography and fluvial systems investigated
- Invasive species

- Impacts and their directionality (negative, positive, or neutral)
- Impacted ecosystem components
- Type of study (e.g. correlational, experimental, etc.)
- Time (did the level of knowledge change over time?)

Additionally, we will delineate potential changes in impact magnitude by species under current climate change scenarios based on the available literature.

Methods

Search string

We will conduct multiple systematic searches, one for each of our focus species. For each search, we will use as keywords the scientific name of a species and “impact”, formatted for Web of Science (WOS). For example:

Elaeagnus angustifolia AND impact*

The selected search string is purposely broad. Searches including keywords associated with the target ecosystem (riparian, foreshore, freshwater, wetland, aquatic, etc.) and geographic area (British Columbia, Canada, North America, etc.) were deemed to be too restrictive. A broader search allows for capturing additional studies that either use different keywords or investigate impacts in different circumstances and yet might be relevant to the target ecosystem.

We tested the comprehensiveness of searches using two pilot species, the Russian Olive (*Elaeagnus angustifolia*) and the Canary Reed Grass (*Phalaris arundinacea*). For each species, we selected 5 primary articles, which used a variety of keywords (e.g. impact, effect, alter, change, consequence, see Additional file 1: Appendix 1 for the full list). Then, we used the search strings to extract studies from WOS and we extracted references from CABI and review studies for pilot species. All studies were detected by search strings. These two species aided the iterative development of the protocol and will be included in the systematic map.

Bibliographic sources

We will conduct searches in WOS, accessing the core database using an institutional licence (University of British Columbia). The core database assigns metadata to a study based exclusively on the information provided by the publisher and journal. Since other databases assign additional metadata to a study, some material might go undetected despite meeting our criteria. We will expand our search to all databases and then refine it to the core collection. This will identify studies that match our keywords across all databases but are only present in the core collection, and thus accessible to the authors (Mathew Vis-Dunbar, UBC librarian, pers. comm. 2023). Additionally, we will screen all references in the CABI

Invasive Species Compendium factsheet for each species, except for references in the *Distribution References* section. Review studies that fit the criteria for inclusion will be used as sources as well, and references extracted and screened.

We will also scope organization websites across North America at different administrative levels. We will assess international (outside Canada), federal (Canada), provincial (British Columbia) and local (regions within British Columbia) organizations. We will search for the focus species name and the word “invasive” in the following organization websites:

- Canadian Weed Society
- British Columbia Inter-Ministry Invasive Species Working Group
- Canadian Council of Invasive Species
- Invasive Species Centre
- Okanagan Basin Waterboard
- North American Invasive Species Management Association (NAISMA)
- The National Environmental Coalition on Invasive Species (NECIS)
- United States Department of Agriculture (USDA)
- National Invasive Species Council
- All local associations in British Columbia (e.g. Boundary Invasive Species Society, East Kootenay Invasive Species Council, Okanagan and Similkameen Invasive Species Society, etc.)

We will conduct the same query in the following searchable catalogues of government documents:

- Canadian Federal Science Library Network
- Legislative library of British Columbia

These sources will allow for capturing also the grey literature. WOS identifies dissertations and conference proceedings, especially if expanding searches to all databases, while the CABI, review papers and organizational websites will identify technical reports.

Accessing multiple databases will help reduce location and index biases (i.e. not all journals are indexed in all databases, incomplete or poor indexing, [45]).

Screening and inclusion criteria

The screening process will include two stages. First, we will screen titles and abstracts. If the information is insufficient to make a decision, we will assess the full manuscript as well. These steps will be applied to all studies, regardless of the source they were extracted from. A single reviewer will conduct the screening (FM). A random subset of studies will also be assessed by a second

reviewer (JP) at both stages (Stage 1 = 5%, Stage 2 = 10%). We will appraise consistency using Cohen’s kappa statistics and set 0.6 as a threshold [60, 61]. If consistency is below the cut-off limit, screening and inclusion criteria will be adjusted for clarity. All disagreements will be discussed and resolved. Any study authored by one of the systematic reviewers that meets the criteria for inclusion will be assessed by the other reviewer at every stage of the process.

We will screen both commercially published and grey literature, but not personal communications or expert opinions. Including grey literature reduces the risk of publication and citation biases (i.e. significant results are more likely to be published and cited than non-significant results, [45, 48]). We will consider only material in English. To minimize language bias (i.e. significant results are more likely to be published in English, [45, 48]), we will assess the title and abstract if translated into English. Studies will be included irrespective of the magnitude, type or directionality of the impact (negative, positive or neutral), and irrespective of the statistical significance of reported results. This will help reduce the prevailing paradigm bias (i.e. a bias towards studies supporting the prevailing paradigm; in this case, invasive species’ impacts are extensive and negative, [26, 45, 48]). The time span includes all studies up to the day the search will be conducted, countering temporal bias (i.e. older studies might be overlooked, [45, 54]). Finally, we will include studies regardless of study design (e.g. experimental, observational, etc.).

We will include studies if they:

- (1) Refer to the non-native invasive plant species searched. We defined as invasive widespread, impactful non-native species.
- (2) Focus on its abiotic and biotic impacts. We defined impacts as measurable changes caused by non-native species on a target ecosystem.
- (3) Investigate such impacts in riparian and foreshore ecosystems. Riparian ecosystems are defined as areas adjacent to streams or rivers (flowing water), while foreshore ecosystems are defined as the land adjacent to still (non-flowing) water bodies.
- (4) Within North America (i.e. Canada and U.S.A.).

We will include all studies in North America because many environmental conditions and invasive species will be shared between British Columbia and other regions within Canada and the U.S. However, including all studies in North America might capture information not relevant to British Columbia. For instance, studies might investigate the impacts of invasive plant species on abiotic and biotic components absent in our study system.

Such cases will be excluded, and exclusions justified. Similarly, we will justify all other exceptions [62].

Study validity assessment

We assessed the validity of each study based only on the eligibility criteria.

Data coding

For each study at the full-text screening stage, we will provide the following information:

1. Bibliographic information
 - a) Authors list
 - b) Article title
 - c) Publication year
 - d) Bibliographic source
2. Inclusion criteria
 - a) Exposure: Focuses on target species (Y/N)
 - b) Exposure: Focuses on abiotic and biotic impacts (Y/N)
 - c) Population: Focuses on riparian and foreshore ecosystems (Y/N)
 - d) Population: Within North America (Y/N)
3. Screening stage
 - a) Excluded at full-text stage
 - b) Included
 - c) Exceptions
4. Additional information
 - a) Duplicate (Y/N)
 - b) Notes

For included studies only, we will provide also the following information:

1. Bibliographic information
 - a) Authors list
 - b) Article title
 - c) Publication year
2. Information on impacts
 - a) Impact description

- b) Ecosystem component impacted (e.g. species, soil, etc.)
- c) Magnitude of impact
- d) Impact direction (negative, positive, neutral)

3. Additional information

- a) Geographic region
- b) Study Design (i.e. field or laboratory experiment, correlation or direct observation)
- c) Notes

We will compile subsection 3c. *Exceptions* on a case-by-case basis. For included studies, we will provide information by impact so that if a study investigated more than one, there will be a number of entries equivalent to the number of impacts assessed.

Meta-data extraction

Studies included in the systematic literature map will undergo a full-manuscript screening to identify the investigated impact (or impacts). We will provide a description of the investigated impacts and the ecosystem component impacted. Then, we will categorize impacts by their magnitude and directionality. Impacts magnitude will be assessed following previous work, modified to include both positive and negative impacts [30–32]:

- Minimal: The impact is unlikely or negligible.
- Minor: It causes changes in the fitness of individuals in the native biota, but no changes in native population densities.
- Moderate: It causes changes in the population densities of native species, but no changes to the structure of communities or the abiotic or biotic components of ecosystems.
- Major: It causes the local or population extinction/introduction of at least one native species, and leads to reversible/transient changes in the structure of communities and the abiotic or biotic components of ecosystems.
- Massive: It leads to the replacement and local extinction/introduction of multiple native species, and produces irreversible changes in the structure of communities and the abiotic or biotic components of ecosystems.

Synthesis and presentation

For each species, we will provide a first database with all studies included at the full-text screening and a reason for exclusions at this stage. A second database with the studies included in the map, along with a graphical

representation of the screening process. Both databases will contain corresponding coded metadata (see *Data Coding* section). We will import studies included in the review into a reference manager and share them as a public library to facilitate accessibility. We will develop a graphical representation of riparian ecosystems, representing identified impacts and their magnitude and directionality for each species. Then, we will create a matrix combining multiple species (as rows) and impacts (as columns) to illustrate the collective impacts of the focus species. Descriptive statistics will be used to answer secondary questions. We will provide the geographic distribution of studies, visualize publication trends over time, and illustrate differences in species and impacts research effort. We will use co-occurrence matrices to identify research effort biases [63]. Lastly, we will provide a narrative synthesis of results for both main and secondary questions. The narrative synthesis will focus on (i) species and impact prioritization, (ii) clusters and gaps in present knowledge, (iii) predicted variations in impact magnitude and direction under current climate change scenarios, and (iv) avenues for future research.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13750-023-00320-3>.

Additional file 1. List of primary studies selected for the comprehensive assessment of the searches.

Additional file 2. ROSES Checklist.

Acknowledgements

We thank Mathew Vis-Dunbar (UBC librarian) for support and guidance in developing search strings and extracting information from bibliographic databases. We thank all the stakeholders for the feedback provided throughout the process.

Author contributions

FM drafted the protocol with input from JP and CM. All authors read and approved the final manuscript.

Funding

FM is funded by the Ministry of Forests, British Columbia, Canada, and by the Irving K. Faculty of Science at the Okanagan campus of UBC. JP acknowledges financial support from the Natural Science and Engineering Council of Canada (Discovery Grant 2020-06543).

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Biology, and Institute for Biodiversity, Resilience, and Ecosystem Services, I.K. Barber Faculty of Science, University of British Columbia Okanagan, 1177 Research Road, Kelowna, BC V1V 1V7, Canada. ²Agriculture and Agri-Food Canada, Summerland Research and Development Centre, 4200 Highway #97, Summerland, BC V0H 1Z0, Canada.

Received: 20 July 2023 Accepted: 14 November 2023

Published online: 14 December 2023

References

- Poff B, Koestner KA, Neary DG, Henderson V. Threats to riparian ecosystems in western north America: an analysis of existing literature. *JAWRA J Am Water Resour Assoc.* 2011;47(6):1241–54.
- Stella JC, Rodríguez-González PM, Dufour S, Bendix J. Riparian vegetation research in Mediterranean-climate regions: common patterns, ecological processes, and considerations for management. *Hydrobiologia.* 2013;719(1):291–315.
- Ragan K, Schipper J, Bateman HL, Hall SJ. Mammal use of riparian corridors in semi-arid Sonora, Mexico. *J Wildl Manag.* 2023;87(1):e22322.
- Koundouri P, Boulton AJ, Detry T, Souliotis I. Ecosystem services, values, and societal perceptions of intermittent rivers and ephemeral streams. In: Detry T, Bonada N, Boulton AJ, editors. *Intermittent rivers and ephemeral streams.* Amsterdam: Elsevier; 2017. p. 455–76.
- Patten DT. Riparian ecosystems of semi-arid North America: diversity and human impacts. *Wetlands.* 1998;18(4):498–512.
- Knopf FL, Johnson RR, Rich T, Samson FB, Szaro RC. Conservation of riparian ecosystems in the United States. *Wilson Bull.* 1988;100(2):272–84.
- Erős T, Kuehne L, Dolezsai A, Sommerwerk N, Wolter C. A systematic review of assessment and conservation management in large floodplain rivers: actions postponed. *Ecol Ind.* 2019;98:453–61.
- Fryirs KA, Zhang N, Duxbury E, Ralph T, Fryirs KA, Zhang N, et al. Rivers up in smoke: impacts of Australia's 2019–2020 megafires on riparian systems. *Int J Wildland Fire.* 2022;31(7):720–7.
- Brierley GJ, Hikuroa D, Fuller IC, Tunnicliffe J, Allen K, Brasington J, et al. Reanimating the strangled rivers of Aotearoa New Zealand. *WIREs Water.* 2023;10(2):e1624.
- Stella JC, Bendix J. Chapter 5—Multiple stressors in riparian ecosystems. In: Sabater S, Elozeigi A, Ludwig R, editors. *Multiple stressors in river ecosystems.* Amsterdam: Elsevier; 2019. p. 81–110.
- Airoldi L, Beck M. Loss, status and trends for coastal marine habitats of Europe. In: Gibson R, Atkinson R, Gordon J, editors. *Oceanography and marine biology: an annual review, vol. 2007*4975. Boca Raton: CRC Press; 2007. p. 345–405.
- Johnson PT, Olden JD, Vander Zanden MJ. Dam invaders: impoundments facilitate biological invasions into freshwaters. *Front Ecol Environ.* 2008;6(7):357–63.
- Chiu MC, Leigh C, Mazor R, Cid N, Resh V. Anthropogenic threats to intermittent rivers and ephemeral streams. In: Detry T, Bonada N, Boulton A, editors. *Intermittent rivers and ephemeral streams.* Amsterdam: Elsevier; 2017. p. 433–54.
- Greene SL. A roadmap for riparian invasion research. *River Res Appl.* 2014;30(5):663–9.
- Post GC, Chang H, Banis D. The spatial relationship between patterns of disappeared streams and residential development in Portland, Oregon, USA. *J Maps.* 2022;18(2):210–8.
- Sweeney BW, Bott TL, Jackson JK, Kaplan LA, Newbold JD, Standley LJ, et al. Riparian deforestation, stream narrowing, and loss of stream ecosystem services. *Proc Natl Acad Sci.* 2004;101(39):14132–7.
- Swift BL. Status of riparian ecosystems in the United States. *JAWRA J Am Water Resour Assoc.* 1984;20(2):223–8.
- Aguiar FCF, Ferreira MT. Plant invasions in the rivers of the Iberian Peninsula, south-western Europe: a review. *Plant Biosyst.* 2013;147(4):1107–19.
- Arianoutsou M, Delipetrou P, Celesti-Grapow L, Basnou C, Bazos I, Kokkoris Y, et al. Comparing naturalized alien plants and recipient habitats

- across an east–west gradient in the Mediterranean Basin. *J Biogeogr.* 2010;37(9):1811–23.
20. Kowarik I, Säumel I. Water dispersal as an additional pathway to invasions by the primarily wind-dispersed tree *Ailanthus altissima*. *Plant Ecol.* 2008;198(2):241–52.
 21. Pysek P, Prach K. Plant invasions and the role of riparian habitats: a comparison of four species alien to Central Europe. *J Biogeogr.* 1993;20(4):413–20.
 22. Barnes WJ. The rapid growth of a population of reed canarygrass (*Phalaris arundinacea* L.) and its impact on some riverbottom herbs. *J Torrey Bot Soc.* 1999;126(2):133–8.
 23. Price EPF, Spyreas G, Matthews JW. Biotic homogenization of regional wetland plant communities within short time-scales in the presence of an aggressive invader. *J Ecol.* 2018;106(3):1180–90.
 24. Bateman HL, Chung-MacCoubrey A, Snell HL. Impact of non-native plant removal on lizards in riparian habitats in the southwestern United States. *Restor Ecol.* 2008;16(1):180–90.
 25. Pendleton R, Pendleton B, Finch D. Displacement of native riparian shrubs by woody exotics: effects on arthropod and pollinator community composition. *Nat Resour Environ Issues.* 2011;16(1):25.
 26. Martinez AE, McDowell PF. Invasive reed canarygrass (*Phalaris arundinacea*) and native vegetation channel roughness. *Invasive Plant Sci Manag.* 2016;9(1):12–21.
 27. Simons SB, Seastedt TR. Decomposition and nitrogen release from foliage of cottonwood (*Populus deltoides*) and Russian-olive (*Elaeagnus angustifolia*) in a riparian ecosystem. *Southwest Nat.* 1999;44(3):256–60.
 28. Mineau MM, Baxter CV, Marcarelli AM. A non-native riparian tree (*Elaeagnus angustifolia*) changes nutrient dynamics in streams. *Ecosystems.* 2011;14(3):353–65.
 29. Jacinthe PA, Bills JS, Tedesco LP. Size, activity and catabolic diversity of the soil microbial biomass in a wetland complex invaded by reed canary grass. *Plant Soil.* 2010;329(1):227–38.
 30. Katsanevakis S, Tempera F, Teixeira H. Mapping the impact of alien species on marine ecosystems: the Mediterranean Sea case study. *Divers Distrib.* 2016;22(6):694–707.
 31. Blackburn TM, Essl F, Evans T, Hulme PE, Jeschke JM, Kühn I, et al. A unified classification of alien species based on the magnitude of their environmental impacts. *PLoS Biol.* 2014;12(5):e1001850.
 32. Vimercati G, Probert AF, Volery L, Bernardo-Madrid R, Bertolino S, Céspedes V, et al. The EICAT+ framework enables classification of positive impacts of alien taxa on native biodiversity. *PLoS Biol.* 2022;20(8):e3001729.
 33. Sax DF, Schlaepfer MA, Olden JD. Identifying key points of disagreement in non-native impacts and valuations. *Trends Ecol Evolut.* 2023. <https://doi.org/10.1016/j.tree.2023.03.004>.
 34. Cardou F, Vellend M. Stealth advocacy in ecology and conservation biology. *Biol Cons.* 2023;280:109968.
 35. Canadian Environmental Assessment Agency. Cumulative effects assessment practitioners guide. Ottawa. 1999.
 36. Council on Environmental Quality. National environmental policy act—cumulative effects.
 37. European Environment Agency. Cumulative Impacts. <https://www.eea.europa.eu/help/glossary/eea-glossary/cumulative-impacts>. Accessed July 2023.
 38. Spaling H, Smit B. Cumulative environmental change: conceptual frameworks, evaluation approaches, and institutional perspectives. *Environ Manage.* 1993;17(5):587–600.
 39. Collier N, Austin BJ, Bradshaw CJA, McMahon CR. Turning pests into profits: introduced buffalo provide multiple benefits to indigenous people of northern Australia. *Hum Ecol.* 2011;39(2):155–64.
 40. Follstad Shah JJ, Harner MJ, Tibbets TM. *Elaeagnus angustifolia* elevates soil inorganic nitrogen pools in riparian ecosystems. *Ecosystems.* 2010;13(1):46–61.
 41. Spyreas G, Wilm BW, Plocher AE, Ketzner DM, Matthews JW, Ellis JL, et al. Biological consequences of invasion by reed canary grass (*Phalaris arundinacea*). *Biol Invasions.* 2010;12(5):1253–67.
 42. Rojas IM, Zedler JB. An invasive exotic grass reduced sedge meadow species richness by half. *Wetlands Ecol Manage.* 2015;23(4):649–63.
 43. Katz GL, Tuttle GM, Denslow MW, Norton AP. Legacy effects of Russian olive (*Elaeagnus angustifolia* L.) in a riparian ecosystem three years post-removal. *Wetlands.* 2020;40(6):1897–907.
 44. Gallego-Tévar B, Grewell BJ, Whitcraft CR, Futrell JC, Bárcenas-Moreno G, Castillo JM. Contrasted impacts of yellow flag iris (*Iris pseudacorus*) on plant diversity in tidal wetlands within its native and invaded distribution ranges. *Diversity.* 2022;14(5):326.
 45. Rodgers VL, Scanga SE, Kolozsvary MB, Garneau DE, Kilgore JS, Anderson LJ, et al. Where is garlic mustard? Understanding the ecological context for invasions of *Alliaria petiolata*. *Bioscience.* 2022;72(6):521–37.
 46. Vanderhoeven S, Dassonville N, Meerts P. Increased topsoil mineral nutrient concentrations under exotic invasive plants in Belgium. *Plant Soil.* 2005;275(1):169–79.
 47. Simberloff D, Von Holle B. Positive interactions of nonindigenous species: invasional meltdown? *Biol Invasions.* 1999;1(1):21–32.
 48. Benoit LK, Askins RA. Impact of the spread of *Phragmites* on the distribution of birds in Connecticut tidal marshes. *Wetlands.* 1999;19(1):194–208.
 49. Brooks RK, Barney JN, Salom SM. The invasive tree, *Ailanthus altissima*, impacts understory nativity, not seedbank nativity. *For Ecol Manage.* 2021;489:119025.
 50. Claeson SM, LeRoy CJ, Barry JR, Kuehn KA. Impacts of invasive riparian knotweed on litter decomposition, aquatic fungi, and macroinvertebrates. *Biol Invasions.* 2014;16(7):1531–44.
 51. Katz GL, Shafroth PB. Biology, ecology and management of *Elaeagnus angustifolia* L. (Russian olive) in western North America. *Wetlands.* 2003;23(4):763–77.
 52. Medina-Villar S, Rodríguez-Echeverría S, Lorenzo P, Alonso A, Pérez-Corona E, Castro-Díez P. Impacts of the alien trees *Ailanthus altissima* (Mill.) Swingle and *Robinia pseudoacacia* L. on soil nutrients and microbial communities. *Soil Biol Biochem.* 2016;96:65–73.
 53. Pullin AS, Frampton GK, Livoreil B, Petrokofsky G. Collaboration for Environmental Evidence. Guidelines and standards for evidence synthesis in environmental management. Version 5.1. 2022. <https://environmentalevidence.org/information-for-authors/>. Accessed July 2023.
 54. Haddaway NR, Bernes C, Jonsson BG, Hedlund K. The benefits of systematic mapping to evidence-based environmental management. *Ambio.* 2016;45(5):613–20.
 55. Haddaway NR, Macura B, Whaley P, Pullin AS. ROSES Reporting standards for systematic evidence syntheses: pro forma, flow-diagram and descriptive summary of the plan and conduct of environmental systematic reviews and systematic maps. *Environ Evid.* 2018;7(1):7.
 56. Palacio FX, Callaghan CT, Cardoso P, Hudgins EJ, Jarzyna MA, Ottaviani G, et al. A protocol for reproducible functional diversity analyses. *Ecography.* 2022;2022(11):e06287.
 57. Bellard C, Thuiller W, Leroy B, Genovesi P, Bakkenes M, Courchamp F. Will climate change promote future invasions? *Glob Chang Biol.* 2013;19(12):3740–8.
 58. Perry LG, Reynolds LV, Shafroth PB. Divergent effects of land-use, propagule pressure, and climate on woody riparian invasion. *Biol Invasions.* 2018;20(11):3271–95.
 59. Poland TM, Patel-Weynand T, Finch DM, Miniati CF, Hayes DC, Lopez VM. Invasive species in forests and rangelands of the United States: a comprehensive science synthesis for the United States Forest Sector. Cham: Springer International Publishing; 2021.
 60. Ridley FA, McGowan PJ, Mair L. The scope and extent of literature that maps threats to species: a systematic map protocol. *Environ Evid.* 2020;9(1):23.
 61. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas.* 1960;20:2746.
 62. O’Dea RE, Lagisz M, Jennions MD, Koricheva J, Noble DWA, Parker TH, et al. Preferred reporting items for systematic reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension. *Biol Rev.* 2021;96(5):1695–722.
 63. Ridley FA, Hickinbotham EJ, Suggitt AJ, McGowan PJK, Mair L. The scope and extent of literature that maps threats to species globally: a systematic map. *Environ Evid.* 2022;11(1):1–26.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.